ELSEVIER International Journal of Mass Spectrometry and Ion Processes 171 (1997) v–vi

International Journal of Mass Spectrometry and Ion Processes

CONTENTS

On the reaction of the ethylene oxide radical cation with neutral ethylene oxide
C.E. Whittle, J. Farr, C. Henrickson, D.S. Wilson, J. Hollis, R.W. Holman (Bowling Green, USA) and M.L. Gross (St. Louis, USA) 1

Single electron capture by B^{2+} ions in Ne, Ar, Kr and Xe
C. Leather, A.R. Lee, A.C.R. Wilkins and A.G. Brenton (Swansea, UK) 7

New operating mode of a quadrupole ion trap in mass spectrometry 4. Simulation studies
P. Perrier, T. Nguema, M. Carette, J. Andre, Y. Zerega, G. Brincourt and R. Catella (Marseille, France) 19

Elimination of the multiple collision effect from a mass-analyzed ion kinetic energy profile in collision-induced dissociation of high mass ions
Y.J. Lee and M.S. Kim (Seoul, South Korea) 31

Infrared and collision-induced fragmentation of iron ethoxide cations
D.A. Kirkwood and A.J. Stace (Brighton, UK) 39

Selected ion flow tube study of the reactions of O^- and O_2^- with $CHCl_2F$, $CHClF_2$, CHF_3, CH_2ClF, CH_2F_2, CH_3F, CHF_2CHF_2, CH_2FCF_3, and CH_3CHF_2
R. Peverall, R.A. Kennedy, C.A. Mayhew (Birmingham, UK) and P. Watts (Salisbury, UK) 51

A peculiar oxygen isotope fractionation in the production of ReO_4^- ion by thermal ionization
Y.Z. Liu, A. Masuda and M. Inoue (Tokyo, Japan) 73

Generation and characterization of ionic and neutral (methylthio)oxophosphane $(CH_3S\text{-}P=O)^{+/o}$ and (methoxy)oxophosphane $(CH_3O\text{-}P=O)^{+/o}$ by neutralization-reionization mass spectrometry
S. Vivekananda and R. Srinivas (Hyderabad, India) 79

Dissociative ionization of silane by electron impact
R. Basner, M. Schmidt, H. Deutsch (Greifswald, Germany), V. Tarnovsky and K. Becker (New York, USA) 83

Fragmentation of $F_3SiCH_2CH_2Si(CH_3)_3$ vapour following Si:2p core-level photoexcitation. A search for a site-specific process in complex molecules
S.-i. Nagaoka, T. Fujibuchia (Matsuyama, Japan), J. Ohshita, M. Ishikawa (Higashi-Hiroshima, Japan) and I. Koyano (Hyogo, Japan) 95

High precision $^{230}Th/^{232}Th$ and $^{234}U/^{238}U$ measurements using energy-filtered ICP magnetic sector multiple collector mass spectrometry
X. Luo, M. Rehkämper, D.-C. Lee and A.N. Halliday (Ann Arbor, USA) 105

Extension of the DM formalism for the calculation of cross sections for the multiple ionization of atoms to the formation of highly charged ions
H. Deutsch (Greifswald, Germany), K. Becker (Hoboken, USA), D.P. Almeida (Santa Catarina, Brazil) and T.D. Märk (Innsbruck, Austria) 119

Generation of an $H_3^+(v=0)$ beam and its diagnosis from the observation of its polar dissociation in collision with He
F.B. Yousif, G. Hinojosa, J. de Urquijo, C. Cisneros and I. Alvarez (Morelos, Mexico) 127

Collision-induced dissociation and post-source decay of model dodecapeptide ions containing lysine and glycine
X. Zhang, J. Jai-nhuknan and C.J. Cassady (Oxford, USA) 135

Electron impact and multiphoton ionization and fragmentation of molybdenum-cyclopentadienyl-dicarbonyl-nitrosyl at 351, 248 and 193 nm
D. Bruch, G. von Bünau (Siegen, Germany) and J. Opitz (Stuttgart, Germany) 147

A triple sector, guided-ion-beam mass spectrometer for cluster ion and fullerene scattering
Y.J. Basir, J.F. Christian, Z. Wan and S.L. Anderson (Salt Lake City, USA) 159

Ionization cross-sections and ionization efficiency curves from polarizability volumes and ionization potentials
P.W. Harland and C. Vallance (Christchurch, New Zealand) 173

Investigations on the ion energy distribution in an analytical radiofrequency glow discharge and its mass spectrometrical application
R. Jäger, J.S. Becker, H.-J. Dietze (Jülich, Germany) and J.A.C. Broekaert (Dusseldorf, Germany) 183

The study of ripple formation and degradation of depth resolution in GaAs and $Al_xGa_{1-x}As$ structures
K. Elst, A. Adriaens and F. Adams (Antwerpen, Belgium) 191
Beam distortions and their compensation in sector field mass spectrometers
M.I. Yavor, A.S. Berdnikov (St. Petersburg, Russia) and H. Wollnik (Giessen, Germany) 203
Dissociation mechanism of electronically excited C_3H_4 isomers by charge inversion mass spectrometry
S. Hayakawa, H. Endoh (Sakai, Japan), K. Arakawa and N. Morishita (Gunma, Japan) 209
Ab initio study of the chloromethane dication fragmentation
D. Duflot (Villeneuve d'Ascq, France) 215
Adsorption in gas mass spectrometry II. Effects on the measurement of isotope amount ratios
R. Gonfiantini, S. Valkiers, P.D.P. Taylor and P. De Bièvre (Geel, Belgium) 231
Ethyl loss kinetics of tetraethylsilane ions studied by TPEPICO
J.W. Keister, P. Tomperi and T. Baer (Chapel Hill, USA) 243
Probing radial and axial secular frequencies in a quadrupole ion trap
M. Carette, P. Perrier, Y. Zerega, G. Brincourt, J.C. Payan and J. Andre (Marseille, France) 253
Calibrated measurements of the isotopic composition and atomic weight of the natural Li isotopic reference material IRMM-016
H.P. Qi, P.D.P. Taylor, M. Berglund and P. De Bièvre (Geel, Belgium) 263
Ab initio calculations on a possible pathway for the formation of C_3H_3 fragment ions from the benzene radical cation
W.J. van der Hart (Leiden, The Netherlands) 269
Electronic structures of $(SO_2)_n^-$ as studied by photoelectron spectroscopy
T. Tsukuda, T. Hirose (Tokyo, Japan) and T. Nagata (Tokyo and Okazaki, Japan) 273

Author Index 281

Subject Index 283

Short communication:
Electron impact ionisation of molecules at high energies
V. Saksena, M.S. Kushwaha (Gwalior, India) and S.P. Khare (Meerut, India) L1

Letters to the Editor:
Solvation of magnesium oxide clusters with water in direct laser vaporization
X.-H. Liu, X.-G. Zhang, X.-Y. Wang and N.-Q. Lou (Dalian, China) L7
Generation and characterization of ionic and neutral thiocarbonylisocyanate $[SCNCO]^{+/}$ by neutralization—reionization mass spectrometry (NRMS)
S. Vivekananda, R. Srinivas (Hyderabad, India) and J.K. Terlouw (Hamilton, Canada) L13